

J/ψ electromagnetic production associated with light hadrons at B factories

Yu-Jie Zhang ^(a,d), Bai-Qing Li ^(b), and Kui-Yong Liu ^(c)

*(a) Key Laboratory of Micro-nano Measurement-Manipulation
and Physics (Ministry of Education) and School of Physics,
Beihang University, Beijing 100191, China*

(b) Department of Physics, Huzhou Teachers College, Huzhou 313000, China

(c) Department of Physics, Liaoning University, Shenyang 110036, China

(d) Kavli Institute for Theoretical Physics China, CAS, Beijing 100190, China

Abstract

The electromagnetic productions of J/ψ associated with light hadrons(LH) and leptonic pairs($\mu^+\mu^-$, $\tau^+\tau^-$) at B factories are studied. We find that the direct electromagnetic production cross sections of $J/\psi(\psi(2S))$ associated with light hadrons is about 0.10(0.04) pb. The direct production cross sections of $J/\psi(\psi(2S))$ associated with $\mu^+\mu^-$ is about 0.056(0.020) pb. If we include the contributions from $\psi(2S)$ decay, we can get the prompt cross section $\sigma[e^+e^- \rightarrow J/\psi + \mu^+\mu^-] = (0.068 \pm 0.002)$ pb, about $(16 \pm 5)\%$ of the Belle data $\sigma[e^+e^- \rightarrow J/\psi + X_{non-c\bar{c}}] = (0.43 \pm 0.09 \pm 0.09)$ pb, meanwhile the $e^+e^- \rightarrow J/\psi + \tau^+\tau^-$ process only contributes 2%. The prompt cross section $\sigma[e^+e^- \rightarrow J/\psi + \text{Light Hadrons}] = (0.121 \pm 0.005)$ pb is about $(28 \pm 8)\%$ of the Belle data.

PACS numbers: 13.66.Bc, 12.38.Bx, 14.40.Pq

I. INTRODUCTION

Heavy quarkonium physics is an important ground to test quantum chromodynamics (QCD) both perturbatively and non-perturbatively. Non-relativistic quantum chromodynamics (NRQCD) factorization approach[1] has achieved a series of successes in describing heavy quarkonium production and annihilation decay. However, there are still some predictions which are less satisfactory. For more details, see a concise review[2]. Among the problematic comparisons with experiment, the large discrepancy between theoretical predictions and experimental data on the charmonium production in e^+e^- annihilation has drawn much attention recently.

In recent years, the B factories has provided systematic measurements on charmonium production. Some results are puzzling, because of the large gap between the measurements and the theoretical predictions. For example, the cross section $\sigma(e^+ + e^- \rightarrow J/\psi + \eta_c)$, measured by Belle Collaboration[3] and Babar Collaboration[4], is almost one order-of-magnitude larger than the leading-order(LO) predictions[5–7]. By introducing the QCD perturbative correction[8, 9], and in combination with relativistic correction[10, 11], this discrepancy was largely resolved. Besides the challenges in the exclusive process, the large ratio of J/ψ production associated with charmed hadrons is also confusing, which was measured by Belle[3]

$$R_{c\bar{c}} = \frac{\sigma[e^+e^- \rightarrow J/\psi + c\bar{c}]}{\sigma[e^+e^- \rightarrow J/\psi + X]} = 0.59_{-0.13}^{+0.15} \pm 0.12. \quad (1)$$

In contrast to the leading-order NRQCD predictions, this ratio is only about 0.1[12–14]. The next-leading-order(NLO) QCD corrections were also introduced in J/ψ inclusive production to resolve the discrepancy between experimental measurements and LO calculations. The NLO QCD corrections to $e^+e^- \rightarrow J/\psi c\bar{c}$ process enhance the cross section with a K factor of about 1.8[15, 16], and only about 20 percent for $e^+e^- \rightarrow J/\psi gg$ process [17, 18]. So the discrepancy is greatly alleviated. To check what role does the color-octet process play, the NLO QCD corrections to color-octet J/ψ inclusive production was calculated[19]. Combining the relativistic corrections to $e^+e^- \rightarrow J/\psi gg$ [20], it may imply that the values of color-octet matrix elements are much smaller than the expected ones which are estimated by using the naive velocity scaling rules.

Another interesting topic is the $e^+e^- \rightarrow J/\psi + X_{non-c\bar{c}}$ production. Most recently, the prompt J/ψ production in association with charmed and non-charmed final particles was

measured[21]

$$\begin{aligned}
\sigma(e^+e^- \rightarrow J/\psi + X) &= (1.17 \pm 0.02 \pm 0.07) pb, \\
\sigma(e^+e^- \rightarrow J/\psi + c\bar{c}) &= (0.74 \pm 0.08^{+0.09}_{-0.08}) pb \\
\sigma(e^+e^- \rightarrow J/\psi + X_{non-c\bar{c}}) &= (0.43 \pm 0.09 \pm 0.09) pb.
\end{aligned} \tag{2}$$

These processes are investigated in Ref.[20, 22], the results show that including both the $O(\alpha_s)$ radiative correction and the $O(v^2)$ relativistic correction, the color-singlet contribution to $e^+e^- \rightarrow J/\psi gg$ has saturated the latest observed cross section $e^+e^- \rightarrow J/\psi + X_{non-c\bar{c}}$ measured by Belle.

Aside from the above QCD process, the pure QED process should also be considered[23]. Especially, in our paper, we calculate virtual-photon-associated production $\sigma(e^+e^- \rightarrow J/\psi\gamma^*) \times B(\gamma^* \rightarrow l\bar{l}(\text{or } Light\ Hadrons))$. The rest of the paper is organized as follows. In Section II, we will give the formulations of $e^+e^- \rightarrow J/\psi + \mu^+\mu^-$. In Section III, the QED production of $e^+e^- \rightarrow J/\psi + LH$ is discussed. In section IV, we will give the numerical results and discussion. Finally we summarize our results in section V.

II. THE FORMULATIONS OF $e^+e^- \rightarrow J/\psi + \mu^+\mu^-$

In NRQCD factorization scheme, the cross section of $e^+e^- \rightarrow J/\psi + \mu^+\mu^-$ can be described as follows

$$\begin{aligned}
&\mathcal{A}(e^+(k_1)e^-(k_2) \rightarrow J/\psi(2p) + \mu^+(p_1) + \mu^-(p_2)) \\
&= \sqrt{C_S} \sum_{L_z S_z} \sum_{s_1 s_2} \sum_{jk} \langle \frac{1}{2}s_1; \frac{1}{2}s_2 | SS_z \rangle \langle LL_z; SS_z | JJ_z \rangle \langle 3j; \bar{3}k | 1 \rangle \times \\
&\quad \mathcal{A}(e^+e^- \rightarrow c_j^{s_1}(p) + \bar{c}_k^{s_2}(p) + \mu^+(p_1) + \mu^-(p_2))
\end{aligned} \tag{3}$$

where $\langle 3j; \bar{3}k | 1 \rangle = 1/\sqrt{3}$, $\langle s_1; s_2 | SS_z \rangle$, $\langle LL_z; SS_z | JJ_z \rangle$ are respectively the color-SU(3), spin-SU(2), and angular momentum Clebsch-Gordan coefficients for $c\bar{c}$ pairs projecting out appropriate bound states. $\mathcal{A}(e^+e^- \rightarrow c_j(p) + \bar{c}_k(p) + \mu^+(p_1) + \mu^-(p_2))$ is the scattering amplitude for $c\bar{c}$ production. The coefficient C_S can be related to the radial wave function of the bound state and reads

$$C_S = \frac{1}{4\pi} |R_S(0)|^2. \tag{4}$$




FIG. 1. The Feynman diagrams of $e^+e^- \rightarrow J/\psi \mu^+ \mu^-$.

We introduce the spin projection operators $P_{SS_z}(p, q)$ as

$$P_{SS_z}(p, q) \equiv \sum_{s_1 s_2} \langle s_1; s_2 | SS_z \rangle v(p - q; s_1) \bar{u}(p + q; s_2). \quad (5)$$

Expanding $P_{SS_z}(p, q)$ in terms of the relative momentum q , we get the projection operators at leading term of q , which will be used in our calculation, as follows

$$P_{1S_z}(p, 0) = \frac{1}{\sqrt{2}} \not{\epsilon}^*(S_z)(\not{p} + M/2). \quad (6)$$

where M is the mass of the charmonium. It is two times of charm quark mass m in the non-relativistic approximation. The polarized cross section can be calculated by defining the longitudinal polarization vector as follows

$$\epsilon_L^\mu(p) = \frac{2p^\mu}{M} - \frac{Mn^\mu}{2n \cdot p}, \quad (7)$$

where $4p^2 = M^2$ and $n^\mu = (1, -\vec{p}/|\vec{p}|)$.

The Feynman diagrams of $e^+e^- \rightarrow J/\psi \mu^+ \mu^-$ are shown in Fig.1. The process of final states with plus charge parity is depicted in diagram Fig.1(a, b) and denoted as $C = +$. The process of final states with minus charge parity is depicted in diagram Fig.1(c, d) and denoted as $C = -$. The $C = +$ process is dominant, and the $C = -$ process is suppressed by a factor of

$$f \sim \ln \left(\frac{s}{4m_\mu^2} \right) \frac{s}{M^2}, \quad (8)$$

here the logarithm term come from quasi-collinear divergence with $m_\mu^2 \ll s$. The s/M^2 term come the photon propagator. One can get $f \sim 10^{-2}$ for $e^+e^- \rightarrow J/\psi \mu^+ \mu^-$ at $\sqrt{s} = 10.6$ GeV for B factories.

III. THE QED PRODUCTION OF $e^+e^- \rightarrow J/\psi + LH$

Similar to the $e^+e^- \rightarrow J/\psi\mu^+\mu^-$ process, the fragment process is also dominant in $e^+e^- \rightarrow J/\psi + \text{Light Hadrons}$ process. The fragment process can be considered as $e^+e^- \rightarrow \gamma^*\gamma^*$, then the virtual photons fragment into J/ψ and light hadrons respectively. By using the approach of the calculation of the hadronic part of the muon $g-2$ [24], this process can be described as

$$\frac{d\sigma^{QED}[e^+e^- \rightarrow J/\psi + LH]}{dm_{LH}^2} \sim \frac{d\sigma[e^+e^- \rightarrow J/\psi + \mu^+\mu^-]}{dm_{\mu^+\mu^-}^2} \times R^{had}(m_{\mu^+\mu^-}^2) \Big|_{m_{\mu^+\mu^-}=m_{LH}}, \quad (9)$$

where

$$R^{had}(\Lambda^2) = \frac{\sigma[e^+e^- \rightarrow \text{hadrons}]}{\sigma[e^+e^- \rightarrow \mu^+\mu^-]} \Big|_{m_{e^+e^-}=\Lambda^2}, \quad (10)$$

because of the contribution of the $C = -$ process is negligible, after subtracting the effect of $\gamma^* \rightarrow c\bar{c}$ from $R^{had}(\Lambda^2)$ by a naive factor $4/3\Theta(\Lambda - M_{c\bar{c}-The})$, we get

$$\begin{aligned} & \frac{d\sigma^{QED}[e^+e^- \rightarrow J/\psi + LH]}{dm_{LH}^2} \\ &= \frac{d\sigma[e^+e^- \rightarrow J/\psi + \mu^+\mu^-]}{dm_{\mu^+\mu^-}^2} \times \left[R^{had}(m_{\mu^+\mu^-}^2) - \frac{4}{3}\Theta(m_{\mu^+\mu^-}^2 - M_{c\bar{c}-The}^2) \right] \Big|_{m_{\mu^+\mu^-}=m_{LH}} \end{aligned} \quad (11)$$

where Θ is step function, $M_{c\bar{c}-The}$ should be correspond to the $c\bar{c}$ threshold of $M_{J/\psi}$, $2M_D$, etc. The uncertainties should be discussed in the next section.

IV. NUMERICAL RESULT

In numerical calculations, the parameters are selected as [25]:

$$\begin{aligned} M_\mu &= 0.1057 GeV, & M_{J/\psi} &= 3.0969 GeV, & \sqrt{s} &= 10.6 GeV, \\ M_\tau &= 1.7768 GeV, & M_{\psi(2S)} &= 3.686 GeV, & \alpha &= 1/132.33 \end{aligned} \quad (12)$$

The table of $R^{had}(\lambda^2)$ have been given in Ref.[24]. We construct an interpolation of $R^{had}(\Lambda^2)$ corresponding to the table with first-order interpolation and setting $R^{had}(\Lambda^2) = 0$ when Λ is less than λ_{min} in the table.

The wave function at the origin can be extracted from the leptonic width $\Gamma(V \rightarrow l^+l^-)$

$$|R_S(0)|^2 = \frac{m_V^2}{4e_Q^2\alpha^2} \Gamma[V \rightarrow e^+e^-]. \quad (13)$$

The leptonic width of charmonium decays into e^+e^- has been given in Ref.[25]

$$\begin{aligned}\Gamma[J/\psi \rightarrow e^+e^-] &= 5.55 \pm 0.14 \text{keV}, \\ \Gamma[\psi(2S) \rightarrow e^+e^-] &= 2.38 \pm 0.04 \text{keV}.\end{aligned}\tag{14}$$

When we calculate the prompt production cross sections of J/ψ , we take into account the feeddown contribution from $\psi(2S)$ by $B[\psi(2S) \rightarrow J/\psi + X] = (57.4 \pm 0.9)\%$ [25] and ignore the contribution of the other charmonium. Then we can get the direct production cross section of $J/\psi(\psi(2S))$ associated with $\tau^+\tau^-$ and $\mu^+\mu^-$ at B factories as

$$\begin{aligned}\sigma^{direct}[e^+e^- \rightarrow J/\psi + \mu^+\mu^-] &= 56 \pm 2 \text{ fb} \\ \sigma^{direct}[e^+e^- \rightarrow J/\psi + \tau^+\tau^-] &= 6.4 \pm 0.2 \text{ fb}\end{aligned}\tag{15}$$

and

$$\begin{aligned}\sigma^{direct}[e^+e^- \rightarrow \psi(2S) + \mu^+\mu^-] &= 20 \pm 1 \text{ fb} \\ \sigma^{direct}[e^+e^- \rightarrow \psi(2S) + \tau^+\tau^-] &= 1.8 \pm 0.1 \text{ fb}\end{aligned}\tag{16}$$

Most of the uncertainties come from the uncertainty of leptonic width in Eq.(14). The others come from the effect of fine structure constant α and higher order QED corrections and so on. The QCD corrections have been taken into account in the leptonic width of $J/\psi(\psi(2S))$.

The cross sections for $C = -$ process is only 1.6%(1.0%) of that for $C = +$ process in $J/\psi(\psi(2S))$ production associated with $\mu^+\mu^-$. And the ratio is about 6.0%(3.9%) in $J/\psi(\psi(2S))$ production associated with $\tau^+\tau^-$. These results are in agreement with the estimation in Eq.(8). So the contribution of $C = -$ process can be ignored in the calculation of electromagnetic $J/\psi(\psi(2S))$ production associated with light hadrons. Finally, we get the direct production cross section of $J/\psi(\psi(2S))$ associated with light hadrons at B factories as

$$\begin{aligned}\sigma_{QED}^{direct}[e^+e^- \rightarrow J/\psi + LH] &= 100 \pm 5 \text{ fb} \\ \sigma_{QED}^{direct}[e^+e^- \rightarrow \psi(2S) + LH] &= 36 \pm 1 \text{ fb}\end{aligned}\tag{17}$$

here we choose $M_{c\bar{c}-The} = 2M_D$ in Eq.(11). If we choose $M_{c\bar{c}-The} = M_{J/\psi}$, there is a difference of -1 fb . So the uncertainties from $M_{c\bar{c}-The}$ can be ignored. Most of the uncertainties come from R^{had} and the leptonic decay width.




FIG. 2. The J/ψ and $\psi(2S)$ energy spectra of direct production processes $e^+e^- \rightarrow V + \mu^+\mu^-$ ($V = J/\psi, \psi(2S)$).

The energy distributions of direct $J/\psi(\psi(2S))$ production from the processes $e^+e^- \rightarrow J/\psi(\psi(2S))\bar{l}l$ and $e^+e^- \rightarrow J/\psi(\psi(2S))LH$ are shown in Fig.2 and Fig.3, respectively. Unfortunately, the endpoint peak was not measured in Ref.[21]. The polarization of $J/\psi(\psi(2S))$ direct production for $e^+e^- \rightarrow V + \mu^+\mu^-$ ($V = J/\psi, \psi(2S)$) process as a function of the energy of J/ψ is shown in Fig.4. The polarization of J/ψ production associated with light hadrons is similar to the results of $e^+e^- \rightarrow V + \mu^+\mu^-$ process. The angular distributions of direct $J/\psi(\psi(2S))$ production from the processes $e^+e^- \rightarrow V + \mu^+\mu^-$ ($V = J/\psi, \psi(2S)$) and $e^+e^- \rightarrow 2\gamma^* \rightarrow V + LH$ ($V = J/\psi, \psi(2S)$) are shown in Fig.5 and Fig.6, respectively.

Now we give the prompt production cross sections of electromagnetic J/ψ production associated with leptonic pairs and light hadrons as

$$\begin{aligned}
 \sigma_{QED}^{prompt}[e^+e^- \rightarrow J/\psi + \mu^+\mu^- + X] &= 68 \pm 2 \text{ fb} \\
 \sigma_{QED}^{prompt}[e^+e^- \rightarrow J/\psi + \tau^+\tau^- + X] &= 7.4 \pm 0.1 \text{ fb} \\
 \sigma_{QED}^{prompt}[e^+e^- \rightarrow J/\psi + LH] &= 121 \pm 5 \text{ fb}.
 \end{aligned} \tag{18}$$




FIG. 3. The J/ψ and $\psi(2S)$ energy spectra of direct production processes $e^+e^- \rightarrow 2\gamma \rightarrow V + LH$ ($V = J/\psi, \psi(2S)$).




FIG. 4. The helicities of J/ψ and $\psi(2S)$ direct production processes $e^+e^- \rightarrow V + \mu^+\mu^-$ ($V = J/\psi, \psi(2S)$) as a function of the energy of J/ψ and $\psi(2S)$.

V. SUMMARY

In summary, the electromagnetic productions of J/ψ associated with light hadrons(LH) and leptonic pairs($\mu^+\mu^-$, $\tau^+\tau^-$) at B factories are studied. We find that the direct electromagnetic production cross sections of $J/\psi(\psi(2S))$ associated with light hadrons is about 0.10(0.04) pb. The direct production cross section of $J/\psi(\psi(2S))$ associated with $\mu^+\mu^-$




FIG. 5. The angular distributions of direct production processes $e^+e^- \rightarrow V + \mu^+\mu^-$ ($V = J/\psi, \psi(2S)$). Here θ is the angle between $J/\psi(\psi(2S))$ momentum and beam.




FIG. 6. The angular distributions of direct production processes $e^+e^- \rightarrow 2\gamma \rightarrow V + LH$ ($V = J/\psi, \psi(2S)$). Here θ is the angle between $J/\psi(\psi(2S))$ and beam.

is about $0.056(0.020)$ pb. If we include the contribution from $\psi(2S)$ decay, we can get the prompt cross section $\sigma[e^+e^- \rightarrow J/\psi + \mu^+\mu^- + X] = (68 \pm 2)$ pb, about $(16 \pm 5)\%$ of the Belle data $\sigma[e^+e^- \rightarrow J/\psi + X_{non-c\bar{c}}] = (0.43 \pm 0.09 \pm 0.09)$ pb, meanwhile the $e^+e^- \rightarrow J/\psi + \tau^+\tau^-$ process only contributes 2%. The prompt cross section $\sigma[e^+e^- \rightarrow J/\psi + \text{Light Hadrons}] = (0.121 \pm 5)$ fb is about $(28 \pm 8)\%$ of the Belle data. Unfortunately, the endpoint peak of energy distribution for J/ψ electromagnetic production associated with leptonic pairs and

light hadrons was not measured in Ref.[21]. The polarization of J/ψ electromagnetic production associated with light hadrons is transversal, while the polarization of J/ψ inclusive production associated with light hadrons from QCD process is longitudinal[22]. We also notify that the charge parity of final states is plus for the QED process calculated in our paper. And it is minus for color singlet process $e^+e^- \rightarrow J/\psi + gg$ and color octet process $e^+e^- \rightarrow J/\psi + g$.

ACKNOWLEDGMENTS

The authors would like to thank Professor K.T.Chao for useful discussion. This work was supported by the National Natural Science Foundation of China (No 10805002, No 10875055), FANEDD2007B18, the Project of Knowledge Innovation Program (PKIP) of Chinese Academy of Sciences, Grant No. KJCX2.YW.W10 and the Education Ministry of Liaoning Province.

-
- [1] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-ibid. D **55**, 5853 (1997)] [arXiv:hep-ph/9407339].
 - [2] G. T. Bodwin, [arXiv:hep-ph/0212203].
 - [3] K. Abe *et al.* [BELLE Collaboration], Phys. Rev. Lett. **89**, 142001 (2002). [arXiv:hep-ex/0205104].
 - [4] B. Aubert *et al.* [BABAR Collaboration] Phys. Rev. D **72**, 031101(2005)
 - [5] E. Braaten and J. Lee, Phys. Rev. D **67**, 054007 (2003) [Erratum-ibid. D **72**, 099901 (2005)] [arXiv:hep-ph/0211085].
 - [6] K. Y. Liu, Z. G. He and K. T. Chao, Phys. Lett. B **557**, 45 (2003) [arXiv:hep-ph/0211181].
K. Y. Liu, Z. G. He and K. T. Chao, Phys. Rev. D **77**, 014002 (2008).
 - [7] K. Hagiwara, E. Kou and C. F. Qiao, Phys. Lett. B **570**, 39 (2003) [arXiv:hep-ph/0305102].
 - [8] Y. J. Zhang, Y. J. Gao and K. T. Chao, Phys. Rev. Lett. **96**, 092001 (2006) [arXiv:hep-ph/0506076].
 - [9] B. Gong and J. X. Wang, Phys. Rev. D **77**, 054028 (2008);Phys. Rev. Lett. **100**, 181803(2008).

- [10] Z. G. He, Y. Fan and K. T. Chao, Phys. Rev. D **75**, 074011 (2007) [arXiv:hep-ph/0702239].
- [11] G. T. Bodwin, J. Lee and C. Yu, Phys. Rev. D **77**, 094018 (2008) [arXiv:0710.0995 [hep-ph]].
- [12] F. Yuan, C. F. Qiao and K. T. Chao, Phys. Rev. D **56**, 321 (1997) [arXiv:hep-ph/9703438].
- [13] P. Cho and A.K. Leibovich, Phys. Rev. D **54**, 6690 (1996); F. Yuan, C.F. Qiao, and K.T. Chao, Phys. Rev. D **56**, 1663 (1997); S. Baek, P. Ko, J. Lee, and H.S. Song, J. Korean Phys. Soc. **33**, 97 (1998); V.V. Kiselev *et al.*, Phys. Lett. B **332**, 411 (1994);
- [14] E. Braaten and Y. Q. Chen, Phys. Rev. Lett. **76**, 730 (1996) [arXiv:hep-ph/9508373].
- [15] Y. J. Zhang and K. T. Chao, Phys. Rev. Lett. **98**, 092003 (2007) [arXiv:hep-ph/0611086].
- [16] B. Gong and J. X. Wang, Phys. Rev. D **80**, 034020 (2009) arXiv:0904.1103 [hep-ph].
- [17] Y. Q. Ma, Y. J. Zhang and K. T. Chao, Phys. Rev. Lett. **102**, 162002 (2009) [arXiv:0812.5106 [hep-ph]].
- [18] B. Gong and J. X. Wang, Phys. Rev. Lett. **102**, 162003 (2009) [arXiv:0901.0117 [hep-ph]].
- [19] Y. J. Zhang, Y. Q. Ma, K. Wang and K. T. Chao, Phys. Rev. D **81**, 034015 (2010)[arXiv:hep-ph/0911.2166].
- [20] Z. G. He, Y. Fan and K. T. Chao, arXiv:0910.3636 [hep-ph].
- [21] P. Pakhlov *et al.* [Belle Collaboration], Phys. Rev. D **79**, 071101 (2009) [arXiv:0901.2775 [hep-ex]].
- [22] Y. Jia, arXiv:0912.5498 [hep-ph].
- [23] C. H. Chang, C. F. Qiao and J. X. Wang, Phys. Rev. D **57**, 4035 (1998).
- [24] V. V. Ezhela, S. B. Lugovsky and O. V. Zenin, arXiv:hep-ph/0312114.
- [25] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).